

CGPG-95/4-2
 gr-qc/9504014
 April 1995

A Modular Invariant Quantum Theory From the Connection Formulation of (2+1)-Gravity on the Torus

Peter Peldán ¹

Center for Gravitational Physics and Geometry
 Physics Department
 The Pennsylvania State University, University Park, PA 16802, USA

Abstract

By choosing an unconventional polarization of the connection phase space in (2+1)-gravity on the torus, a modular invariant quantum theory is constructed. Unitary equivalence to the ADM-quantization is shown.
 PACS numbers: 04.60.-m, 04.60.Ds, 04.60.Kz

It has recently been shown [1] that there are severe obstructions to constructing a quantum theory for (2+1)-dimensional gravity on a manifold $\mathcal{R} \times T^2$ in the connection representation if we treat all diffeomorphisms as gauge. That is, we require physical states in our Hilbert space to be invariant – or at most transform according to a one-dimensional unitary representation of the diffeomorphism group – under all diffeomorphisms. In [1] only the space-like sector [2, 3, 4] was considered, but the result can easily be transferred to the time-like sector as well.

The purpose of this Letter is to show that this problem can be solved straightforwardly by choosing a different polarization of the phase space [5]. The polarization chosen here leads automatically to the Heisenberg picture of the ADM-formulation [6, 7].

The reason why we get such severe problems in trying to construct a quantum theory in the connection representation with all diffeomorphisms treated as gauge, is that the group of large diffeomorphisms has a very bad behavior on the reduced configuration space; the group of large diffeomorphisms on the torus does not act properly discontinuous in the space of gauge equivalent classes of flat $SO(1, 2)$ connections on the torus. However, we know that the action of this group is nicely behaved in the phase space – except for a region of measure zero – and we should therefore be able to quantize this formulation with the use of a different polarization.

¹Email address: peldan@phys.psu.edu

In the space-like sector of (2+1)-gravity on the torus, we have, after solving all constraints and gauge fixing all local symmetries, the following theory [2, 3, 4]: the reduced phase space is the cotangent bundle over the reduced configuration space $\mathcal{Q} = (\mathcal{R}^2 - \{0, 0\})/\mathcal{Z}_2$ coordinatized by the coordinates μ and λ , where $\mathcal{Z}_2$ acts in $\mathcal{R}^2$ as $\{\mu, \lambda\} \rightarrow \{-\mu, -\lambda\}$. The canonically conjugate momenta are denoted a and b and we have the two fundamental Poisson brackets: $\{\mu, a\} = \frac{1}{2}$ and $\{\lambda, b\} = -\frac{1}{2}$. Note that the $\mathcal{Z}_2$ identification in the phase space reads; $(\mu, \lambda, a, b) \sim -(\mu, \lambda, a, b)$. The Hamiltonian is identically zero. Since all local symmetries have been taken care of, the only remaining gauge redundancies are the large diffeomorphisms and the large $O(1, 2)$ transformations. The large diffeomorphisms on the torus form a group – the modular group, $PSL(2, \mathcal{Z}) = SL(2, \mathcal{Z})/\{\mathbf{1}, -\mathbf{1}\}$ – which is generated by

$$S : \quad (a, \lambda) \rightarrow (b, \mu), \quad (b, \mu) \rightarrow (-a, -\lambda) \quad (1)$$

$$T : \quad (a, \lambda) \rightarrow (a, \lambda), \quad (b, \mu) \rightarrow (b + a, \mu + \lambda) \quad (2)$$

while the remaining large $O(1, 2)$ part is isomorphic to $\mathcal{Z}_2$ and generated by

$$O : \quad (a, \lambda) \rightarrow (a, -\lambda), \quad (b, \mu) \rightarrow (b, -\mu) \quad (3)$$

To quantize this system, I will follow the algebraic quantization program outlined in [8] while treating the above large symmetries as gauge. Alternatively, one may see the quantization below as a straightforward application of standard geometrical quantization [5] with a special polarization of the phase space. Instead of quantizing the fundamental Poisson-algebra in the vector space of complex valued functions in $\mathcal{Q}$, I choose to quantize the Poisson-algebra of the phase space functions²

$$m := \frac{(b + i\mu)}{(a + i\lambda)} \quad \bar{m} := \frac{(b - i\mu)}{(a - i\lambda)} \quad (4)$$

$$p := -i(a - i\lambda)^2 \quad \bar{p} := i(a + i\lambda)^2 \quad (5)$$

in the vector space of complex valued functions of m and $\bar{m}$. Note that these functions automatically takes care of the redundancy identification $(\mu, \lambda, a, b) \sim -(\mu, \lambda, a, b)$, while the large diffeomorphisms and $O(1, 2)$ transformations become

$$S : \quad m \rightarrow -\frac{1}{m}, \quad p \rightarrow m^2 p \quad (6)$$

$$T : \quad m \rightarrow m + 1, \quad p \rightarrow p \quad (7)$$

$$O : \quad m \rightarrow \bar{m}, \quad p \rightarrow -\bar{p} \quad (8)$$

and similarly for their complex conjugate functions. Explicitly, the classical Poisson-algebra is

$$\{m, \bar{m}\} = \{m, p\} = \{\bar{m}, \bar{p}\} = \{p, \bar{p}\} = 0 \quad (9)$$

$$\{m, \bar{p}\} = \{\bar{m}, p\} = -2 \quad (10)$$

²The reasons for this choice is partly that these functions are good coordinates on phase space – the Jacobian is everywhere non zero – and partly that the modular group now has a nice action in $m, \bar{m}$ space except for the region $m - \bar{m} = 0 \Leftrightarrow a\mu - b\lambda = 0$.

which is easily quantized by the representation

$$\hat{m} = m, \quad \hat{\bar{m}} = \bar{m}, \quad \hat{p} = 2i\frac{\partial}{\partial \bar{m}} + \frac{1}{m_2}, \quad \hat{\bar{p}} := 2i\frac{\partial}{\partial m} - \frac{1}{m_2} \quad (11)$$

where $m = m_1 + im_2$. (The representation of $\hat{p}$ and $\hat{\bar{p}}$ is chosen such that formally $\hat{p}^\dagger = \hat{\bar{p}}$ w.r.t the inner-product defined below.)

Next we need to find the physical states, *i.e.* complex valued functions of m and $\bar{m}$ that are invariant under the remaining "gauge transformations" (6), (7) and (8). Since we know that the quotient space $\mathcal{C}/PSL(2, \mathcal{Z})/\mathcal{Z}_2$ simply is the torus moduli space³, $\mathcal{M}$ [9], we automatically know that our physical states are given by functions on this space. Then, we need an inner-product, and since there exist a unique – up to scaling – modular invariant measure [9] on the moduli space, the inner product is naturally chosen to be

$$\langle \Psi_1 | \Psi_2 \rangle := \int_{\mathcal{M}} \frac{dm_1 dm_2}{m_2^2} \bar{\Psi}_1 \Psi_2 \quad (12)$$

Explicitly, the moduli space is $\mathcal{M}$: $|m| \geq 1$, $m_2 > 0$ and $|m_1| \leq \frac{1}{2}$.

To complete the quantization one should now find a complete set of $PSL(2, \mathcal{Z}) \otimes \mathcal{Z}_2$ invariant physical observables, check that they are self adjoint in the Hilbert space, and check if their algebra is irreducibly represented in the Hilbert space. I will not address these issues here.

Thus, this completes the construction of a non-trivial Hilbert space carrying a one-dimensional unitary representation of the modular group.

Now, how is this quantum theory related to the ADM-quantization? Just by comparing the definitions (4)-(5) to the time dependent canonical transformation between half the phase space⁴ of the connection formulation and the ADM-formulation [2], one immediately sees that the functions m , $\bar{m}$, p and $\bar{p}$ are nothing more than the ADM-variables at a specific instant of ADM-time. Thus, the above quantum theory is simply the Heisenberg picture of the ADM Schrödinger picture, and unitary equivalence follows directly.

Finally we may note that although we have managed to construct a non trivial modular invariant theory starting from the connection formulation, our quantum theory is not a connection representation [3, 10]. This immediately implies that we do not have any obvious way to relate this quantum theory to the so called loop-representation [3, 10]. Furthermore, it seems very unlikely that the above construction should have any chance of working in the time-like sector where the modular group does not act properly discontinuous even in the full phase space. Thus the construction of a non trivial modular invariant connection representation still remains as an open problem.

Acknowledgements

I thank Abhay Ashtekar for suggesting solving the problem of ref.[1] by choosing a different polarization. I also thank Don Marolf for discussions.

³Actually, to be completely honest, this is only true for the complex plane minus the real axis. However, since the real axis is a region of measure zero in the complex plane, I will simply neglect it.

⁴Explicitly we have $\Gamma_{ADM} \cong \Gamma_{conn}/\mathcal{Z}_2$, where Γ denote the phase space and $\mathcal{Z}_2$ acts in Γ_{conn} as in eq. (3).

References

- [1] P. Peldán, "Large Diffeomorphisms in (2+1)-quantum Gravity on the Torus", Penn State preprint 95/1-1, gr-qc/9501020
- [2] S. Carlip, "(2+1)-dimensional Chern-Simons gravity as a Dirac square root", Phys. Rev. **D 45** (1992) 3584
S. Carlip, "Observables, Gauge Invariance and Time in (2+1)-Dimensional Quantum Gravity", Phys. Rev. **D42** (1990) 2647
- [3] A. Ashtekar, V. Husain, C. Rovelli, J. Samuel and L. Smolin, "2+1 quantum gravity as a toy model for the 3+1 theory", Class. Quant. Grav. **6** (1989) L185
A. Ashtekar and R. Loll, "New Loop Representations for 2+1 Gravity", Class. Quant. Grav. **11** (1994) 2417, gr-qc/9405031
- [4] J. Louko and D. M. Marolf, "Solution Space of 2+1 Gravity on $\mathcal{R} \times T^2$ in Witten Connection Formulation", Class. Quant. Grav. **11** (1994) 311, gr-qc/9308018
- [5] N. Woodhouse, "Geometric Quantization", (Clarendon Press, Oxford, 1980)
A. Ashtekar and invited contributors, "New Perspectives in Canonical Gravity", (Lecture Notes),(Bibliopolis, Napoli, Italy, 1988)
- [6] A. Hosoya and K. Nakao, "(2+1)-Dimensional Pure Gravity for an Arbitrary Closed Initial Surface", Class. Quant. Grav. **7** (1990) 163
Y. Fujiwara and J. Soda, "Teichmüller Motion of (2+1)-Dimensional Gravity with the Cosmological Constant", Progr. Theor. Phys. **83** (1990) 733
V. Moncrief, "How Solvable is (2+1)-Dimensional Einstein Gravity", J. Math. Phys. **31** (1990) 2978,
"Reduction of the Einstein Equations in 2+1 Dimensions to a Hamiltonian System over Teichmüller Space", J. Math. Phys. **30** (1989) 2907
A. Hosoya and K. Nakao, "(2+1)-Dimensional Quantum Gravity", Prog. Theor. Phys. **84** (1990) 739
- [7] S. Carlip, "Six Ways to Quantize (2+1)-dimensional Gravity", gr-qc/9305020
- [8] A. Ashtekar and R. S. Tate, "An Algebraic Extension of Dirac Quantization: Examples", J. Math. Phys. **35** (1994) 6434, gr-qc/9405073
- [9] T. Miyake, "Modular Forms", (Springer-Verlag, Berlin Heidelberg, 1989)
H. Maass, Lecture Notes: "Lectures on Modular Functions of One Complex Variable", (Tata Institute of Fundamental Research, Bombay, 1964)
A. Terras, "Harmonic Analysis on Symmetric Spaces and Applications I", (Springer Verlag, New York, 1985)
- [10] D. M. Marolf, "Loop Representations for 2+1 Gravity on a Torus", Class. Quant. Grav. **10** (1993) 2625, gr-qc/9303019